Draft version October 5, 2012
Preprint typeset using LaTeX style emulateapj v. 04/21/05

arXiv:1210.1494v1 [astro-ph.HE] 4 Oct 2012

LOW-FREQUENCY QPO FROM THE 11 HZ ACCRETING PULSAR IN TERZAN 5: NOT FRAME DRAGGING

D. Altamirano[1], A. Ingram[2], M. van der Klis[1], R. Wijnands[1], M. Linares[3] & J. Homan[3]

Draft version October 5, 2012

ABSTRACT

We report on 6 RXTE observations taken during the 2010 outburst of the 11 Hz accreting pulsar IGR J17480–2446 located in the globular cluster Terzan 5. During these observations we find power spectra which resemble those seen in Z-type high-luminosity neutron star low-mass X-ray binaries, with a quasi-periodic oscillation (QPO) in the 35–50 Hz range simultaneous with a kHz QPO and broad band noise. Using well known frequency-frequency correlations, we identify the 35–50 Hz QPOs as the horizontal branch oscillations (HBO), which were previously suggested to be due to Lense-Thirring precession. As IGR J17480–2446 spins more than an order of magnitude more slowly than any of the other neutron stars where these QPOs were found, this QPO can not be explained by frame dragging. By extension, this casts doubt on the Lense–Thirring precession model for other low-frequency QPOs in neutron-star and perhaps even black-hole systems.

Subject headings: X-rays: binaries — binaries: close — stars: individual (IGR J17480–2446, Terzan 5)

1. INTRODUCTION

One of the strongest motivations for studying low-mass X-ray binaries (LMXBs) has been the aim to use these systems as probes of fundamental physics. A possible tool for this is provided by the quasi-periodic oscillations (QPOs) in the X-ray light curves of LMXBs. These QPOs have now been observed in many LMXBs containing either neutron stars (NSs) or black holes (BHs), and are usually detected with characteristic frequencies between ∼1 mHz and ∼1 kHz. The QPO frequency, coherence and amplitude usually correlate with the source spectral states and/or X-ray luminosity, observables probably set by the geometry and dynamics of the accretion flow (see, e.g., van der Klis 2006, for a review), supporting the idea that QPOs can be used as probes of the flow of matter in strong-field gravity and hence of fundamental physics.

Persistent NS systems have been generally divided into low-luminosity (“4U”) and high-luminosity (“GX”) atoll sources, and the always high-luminosity Z-sources, based not only on luminosity, but also on the tracks they trace out in color-color (CCD) and hardness-intensity diagrams (HID) and on their correlated rapid X-ray variability (Hasinger & van der Klis 1989). There is no similar subdivision for BH systems.

A number of QPOs and broad-band variability components are often present simultaneously in the power spectra of the X-ray light curves of these systems. These power spectra can be fully described with a phenomenological model comprising several Lorentzian components, where each component is generally labeled by its characteristic frequency (e.g. Belloni et al. 2002). Recently, more physically-motivated models based on Lorentzian mass accretion rate fluctuation spectra in the disk have begun to be explored (see Ingram et al. 2009; Ingram & Done 2010, 2011, 2012). When the multi-Lorentzian model is used, each component is called L_i, and its characteristic frequency is ν_i, where i is an identifying symbol (e.g. Altamirano et al. 2008).

Power spectral features in atoll sources include broad components known as the break component (L_b), the hump L_h, the $L_{\ell ow}$ and narrower components, or QPOs, such as the low-frequency QPO (L_{LF}) and the upper and lower kHz QPOs (L_u and L_ℓ, respectively). For Z-sources, low-frequency QPOs have been labeled differently: the horizontal-branch, normal-branch and flaring-branch oscillations (HBOs, NBOs and FBOs, respectively) depending on which spectral state (or “branch” in the HID) of the source they are most prominent. In Figure 1 we show examples; for a more detailed description, we refer the reader to the review by van der Klis (2006). In addition to the break component (and some other broad-band components like $L_{\ell ow}$), BHs generally show three main types of low-frequency QPOs (Types A, B and C, e.g., Casella et al. 2005) and, in a few cases, high-frequency QPOs (with frequencies between 70 Hz and 450 Hz, e.g., Remillard & McClintock 2006; Belloni et al. 2012).

Similarities in the morphology of their power spectra suggest that same variability components are present in both NS and BHs (e.g., Miyamoto et al. 1993; van der Klis 1994; Olive et al. 1998; Belloni et al. 2002; Linares et al. 2007). Correlations between the frequencies of some of these power-spectral components, and similarities in their characteristics confirm this suggestion, providing intriguing links between NSs and BHs. In particular, it has often been suggested that the HBOs in Z-sources are similar to the LF QPO and hump component in atoll sources (e.g. van Straaten et al. 2003) and in BHs (e.g. Casella et al. 2005).

Studies have been done on the relation between the frequencies of various variability components in compact objects. The two best known frequency-frequency

---

[1] Email: d.altamirano@uva.nl ; Astronomical Institute, “Anton Pannekoek”, University of Amsterdam, Science Park 904, 1098XH, Amsterdam, The Netherlands.
[2] Department of Physics, University of Durham, South Road, Durham DH1 3LE, UK.
[3] Massachusetts Institute of Technology - Kavli Institute for Astrophysics and Space Research, Cambridge, MA 02139, USA.

correlations which involve both NSs and BHs are those known as WK (Wijnands & van der Klis 1999) and PBK (Psaltis et al. 1999a) relations. The WK relation links the frequencies of the well identified L_b noise component in atoll sources and BHs, to the L_h/L_{LF} LF-QPO in atoll and BH sources[4]. Wijnands & van der Klis (1999) showed that the frequencies are well correlated over 3 orders of magnitude. In addition, these authors found that the HBOs in Z-sources also show the same relation with L_b, albeit slightly above the atoll/BH track. The PBK relation links the second highest frequency observed in NS (L_ℓ in the high-luminosity soft state and $L_{\ell ow}$ in the low-luminosity hard state) and BHs ($L_{\ell ow}$ in the low-hard state) on one hand, and low-frequency QPOs on the other (e.g., Belloni et al. 2002). This relation spans nearly three decades in frequency (and even a larger range if one considers QPOs in white dwarfs systems, see, e.g., Warner & Woudt 2002). However, as Psaltis et al. (1999a) note, although the correlation is very suggestive, the relation between components is not conclusive as it combines features with different coherence and amplitudes from different sources.

The WK and PBK relations suggest that physically similar (or identical) phenomena set the frequencies observed in BHs and low- and high-luminosity NSs. If true, then the mechanism that sets their frequency must arise in the accretion disk, i.e., it cannot depend on a solid surface (e.g., Wagoner et al. 2001; Rezzolla et al. 2003).

1.1. *Relativistic orbital motion and the identification of Lense-Thirring precession*

Orbits tilted relative to the equatorial plane of a central spinning object show relativistic nodal precession due to frame dragging. In the Lense-Thirring (LT) approximation (weak field and low spin, see Lense & Thirring 1918) the precession frequency is given by $\nu_{\rm LT} = GJ/\pi c^2 r^3$ where J is the angular momentum of the central object and r the orbital radius.

In this approximation the LT precession frequency around a spinning NS can be written as

$$\nu_{\rm LT} = 13.2{\rm Hz}\ I_{45}\left(\frac{M}{M_\odot}\right)^{-1}\left(\frac{\nu_\phi}{1000{\rm Hz}}\right)^2\left(\frac{\nu_{\rm spin}}{300{\rm Hz}}\right) \quad (1)$$

where ν_ϕ is the orbital frequency, I_{45} the moment of inertia of the NS in units of 10^{45} g cm^2, M its mass, and $\nu_{\rm spin}$ its spin frequency (Stella & Vietri 1998).

Stella & Vietri (1998) proposed that the low-frequency QPOs (HBOs and given WK and PBK, also the L_{LF} and/or L_h) in NS systems represent LT precession of the orbit whose general-relativistic orbital frequency ν_ϕ is given by the upper kHz QPO frequency ν_u. Furthermore, Stella et al. (1999) proposed that via the correlations identified by PBK this interpretation can be extended to LF QPOs seen in BH systems. For the NS case, the predicted and observed frequencies do not match directly: as Stella & Vietri (1998) note, for reasonable values of $I_{45}/(M/M_\odot)$ and $\nu_{\rm spin}$, the predicted precession frequencies were a factor of $\sim$2 lower than the observed ones (or even a larger factor, e.g. Jonker et al. 1998; Morsink & Stella 1999; Psaltis et al. 1999b; Jonker et al. 2000). This discrepancy was then interpreted by assuming that the modulation can be produced at twice the LT precession frequency (see discussion in Stella & Vietri 1998), which is not unreasonable given the bilateral symmetry inherent in the geometry of a tilted precessing orbit.

Based on the works of Liu & Melia (2002) and Fragile et al. (2007), Ingram & Done (2010) recently were able to naturally explain the factor of $\sim$2 discrepancy by relaxing the test-particle assumption, and considering LT precession of a geometrically thick inner accretion flow, where a thin truncated outer disk remains stationary but the inner flow precesses as a solid body.

One of the ingredients in all the above works is that the spin frequency of the NS is in the range 200–600 Hz. Most of the $\nu_{\rm spin}$ measurements in NS-LMXBs are in that range (Patruno & Watts 2012). The recent discovery of a new X-ray transient in the globular cluster Terzan 5 containing an 11 Hz pulsar allows us to examine the frame dragging model in a new regime.

1.2. *Testing LT precession models with IGR J17480–2446*

IGR J17480–2446 was discovered in the globular cluster Terzan 5 on October 10th, 2010, with INTEGRAL. Initial RXTE observations revealed an 11 Hz pulsar (Strohmayer & Markwardt 2010) in a 21.3 hr orbital period binary (Papitto et al. 2011). This spin is slow compared to the 185–650 Hz known spin of the low-magnetic ($\sim$10^8 Gauss) field NS-LMXBs, yet (much) faster than that of the ($5\times10^{-5}-2$) Hz pulsars, which exhibit various characteristics indicating a strong ($\gtrsim 10^{11-12}$ Gauss) magnetic field strength (e.g., Patruno & Watts 2012, for a review). The magnetic field of IGR J17480–2446 may be intermediate in strength ($\sim$10^{10} Gauss, e.g., Cavecchi et al. 2011, Papitto et al. 2011 and Patruno et al. 2012).

Near the outburst peak, at about half the Eddington luminosity (e.g. Chakraborty et al. 2011; Linares et al. 2011), IGR J17480–2446 showed X-ray spectral and variability behavior typical of Z sources (Altamirano et al. 2010), with simultaneous broad-band noise, a QPO at $\sim$48 Hz, and a kHz QPO at $\sim$815 Hz. These power-spectral components resemble those seen in other NS systems, where the $\sim$48 Hz QPOs are the ones identified with LT precession.

2. OBSERVATIONS AND DATA ANALYSIS

We use data from the RXTE Proportional Counter Array (PCA; for instrument information see Jahoda et al. 2006). There were 48 pointed observations, each consisting of a fraction of one to several entire satellite orbits. For the timing analysis we used the Event mode E_125us_64M_0_1s. Leahy-normalized power density spectra (Leahy et al. 1983) were constructed using data segments of 128 seconds and 1/8192 s time bins. All frequency bins between 11.02 and 11.07 Hz were removed to get rid of the pulsar spike. No background or deadtime corrections were performed prior to the calculation of the power spectra. We averaged the power spectra per orbit and per observation, subtracted a modeled Poisson noise spectrum (Zhang et al. 1995) and converted the resulting power spectra to squared fractional rms (van der Klis

---

[4] As noted by Belloni et al. (2002), Klein-Wolt & van der Klis (2008) and van Straaten et al. (2003), L_{LF} and L_h are often close in frequency, separated by no more than a few Hz.



Fig. 1.— *Top:* Representative power spectrum of IGR J17480–2446 (ObsID:95437-01-09-00) during which we detect a low-frequency and a single kHz QPO. *Middle* and *Bottom* panels show power spectra for the Z-source GX17+2 (ObsID:20053-03-02-010) and the atoll source 4U1728–34 (ObsID:20083-01-04-00), respectively. All were made using 2–60 keV RXTE data. Power-spectral components are labeled in each panel.

1995).

For a detailed analysis of the power and energy spectral evolution along the outburst, we refer the reader to Altamirano et al. (2012) and Barret (2012). In the current work, we concentrate on the 6 observations where the low-frequency QPOs and the kHz QPOs were detected.

3. RESULTS

We found LF QPOs with frequencies between ∼35 Hz and ∼50 Hz in 6 observations (9 independent satellite orbits). This QPO was not always present during the whole observation. kHz QPOs with frequencies between ∼800 Hz and 920 Hz (Altamirano et al. 2012; Barret 2012) were detected in 5 observations, simultaneously with the low-frequency QPOs (Altamirano et al. 2012). In the top panel of Figure 1 we show a representative power spectrum of IGR J17480–2446 where we detect a broad feature (zero-centered Lorentzian) with a characteristic frequency of 14.8 ± 0.8 Hz, a QPO at 45.0 ± 0.7 Hz and a kHz QPO at 851 ± 4 Hz. The fractional amplitude of the 45 Hz QPO increases with energy, from undetected in the 2–5 keV (with a 3σ upper limit of 2.9% rms) to 7.3 ± 1.3% rms at ∼20 keV. The statistics are not sufficient to study the QPO phase-lags. The overall power spectral shape resembles that observed in some Z sources, and to a lesser extent, those observed in atoll sources (middle and bottom panels in Figure 1, respectively; see also Altamirano et al. 2012). The frequency range, the quality factor and rms amplitude of the 35-50 Hz QPOs, together with the overall power-spectral shape and the position where the QPOs occur in the HID (Altamirano et al. 2012), are all consistent with these QPOs being the HBO in Z sources. The kHz QPO can be identified as either the upper or the lower kHz QPO (Altamirano et al. 2012; Barret 2012).

We further tested our identifications using the WK relation, as the break component L_b and the low-frequency QPO are easy to identify. In Figure 2 we plot the data from Wijnands & van der Klis (1999). Our points for IGR J17480–2446 are on the main relation, hugging the upper envelope of the atoll sources (black squares), but still below the Z-sources (red triangles). (Slight differences in how ν_b is measured will affect the values by < 15% (Belloni et al. 2002) and hence are immaterial to our conclusions.)

As discussed by Méndez & Belloni (2007), the available results on NS kHz QPOs are inconclusive on whether $\Delta\nu = \nu_u - \nu_\ell$ is related to the spin frequency of the NS as $\Delta\nu = \nu_{\rm spin}$, or as $\Delta\nu = \nu_{\rm spin}/2$, or it is independent of $\nu_{\rm spin}$ and close on average to ∼300 Hz. In Figure 3 we compare IGR J17480–2446's QPO frequencies with those of the HBO and upper kHz QPO in other Z sources (see Altamirano et al. 2012). The data are suggestive of the correct identification of the different components assuming that either the kHz QPO we detect is the upper, or it is the lower and $\Delta\nu \ll$300 Hz. The PBK relation (not shown) uses the frequency of the lower kHz QPO. In this case, the data of IGR J17480–2446 are also on the main correlation if we assume that we are detecting the upper kHz QPO, and that $\Delta\nu$ is around 300 Hz.

4. DISCUSSION



FIG. 2.— WK relation after Wijnands & van der Klis (1999). Black squares, red triangles and blue diamonds are data from atoll sources, Z sources, and BHs, respectively. Green circles are from IGR J17480–2446. Inset shows a zoom-in. No errors are shown in the main figure for clarity.



FIG. 3.— HBO vs. upper kHz QPO frequency for different Z sources (data from Altamirano et al. 2012). Red circles represent our simultaneous detections of the HBO and kHz QPO assuming that the single kHz QPO is the upper one, or that $\Delta\nu$ is of the order of the spin frequency of IGR J17480–2446 (i.e. 11 Hz). For clarity, we show only error bars for IGR J17480–2446 data.



FIG. 4.— General-relativistic orbital frequency ν_ϕ, periastron precession frequency $\nu_{\rm peri}$ and nodal precession frequency $\nu_{\rm nodal}$ as a function of orbital radius r for an 11 Hz pulsar calculated from the expressions in Merloni et al. (1999). Horizontal shaded bands at 35-50 Hz and 800-900 Hz indicate the observed HBO and kHz QPO frequencies. Black shaded area at bottom marks $\nu_{\rm nodal}$ for prograde orbits and $R = 6R_g$; upper and lower bounds are practically independent of mass and correspond to $\beta = I/(MR^2)$ equal 2/3 and 1/6. For $M = 1.4M_\odot$ these bounds correspond to $I_{45}/(M/M_\odot)$ $\sim$2 and $\sim$0.5, respectively. The curve with arrows is an upper limit to $\nu_{\rm nodal}$ for extreme assumptions maximizing this frequency: retrograde orbital motion, $\beta = 2/3$, and NS radius equal to the orbital radius ($R = r$). Even with extreme assumptions theoretical nodal precession frequencies remain far below the observed HBO frequencies. At this low spin ν_ϕ and $\nu_{\rm peri}$ (both for prograde orbits) are practically independent of R and β but they do depend on mass; curves are for $1.4M_\odot$. These curves can match the observed frequencies, but not at the same orbital radius.

In this paper we report on 6 observations of the 11 Hz accreting pulsar IGR J17480–2446 in which we find power spectra which resemble those seen in Z-type high-luminosity NS-LMXBs. The low-frequency QPOs are in the 35-50 Hz range, while the kHz QPOs are in the 800–920 Hz range. Based on the power-spectral characteristics, where the QPOs occur in the HID (Altamirano et al. 2012), and comparison with frequency-frequency correlations, we identify the low-frequency QPO as the HBO. Our results suggest that the kHz QPO is the upper one, although the results are not conclusive. For a more detailed discussion about the identification of the kHz QPO, we refer the reader to Altamirano et al. (2012). In this letter, we discuss our results mainly in the context of Lense-Thirring precession. Comparison of our results with other proposed models for low-frequency QPOs will be reported elsewhere.

4.1. *Lense-Thirring precession*

The LT precession frequency for a test particle as introduced in Section 1.1 depends on the NS spin, its moment of inertia and mass, and on the orbital frequency. For IGR J17480–2446 $\nu_{\rm spin} = 11$ Hz. Depending on whether the NS equation of state (EoS) is soft or stiff, respectively, $I_{45}/(M/M_\odot)$ could be between 0.5 and 2 (Stella & Vietri 1998). Assuming $I_{45}/(M/M_\odot) < 2$, we obtain $\nu_{\rm LT} \lesssim 0.97\,{\rm Hz}\times(\nu_\phi/1000\,{\rm Hz})^2$. If we associate the 800–920 Hz kHz QPO frequency with ν_ϕ, then $\nu_{\rm LT} < 0.82$ Hz, i.e. much lower than the 35-50 Hz QPO we observe.

We can take a step further and estimate the relativistic nodal precession frequency independently of the nature of the kHz QPOs. The LT precession frequency (Section 1.1) can be written as $\nu_{\rm LT} = 2G\nu_{\rm spin}I/c^2r^3$. Writing the NS moment of inertia as $I = \beta\ MR^2$ where β is a dimensionless constant, we have $\nu_{\rm LT} = 2\beta R_g R^2\nu_{\rm spin}/r^3$, where the gravitational radius $R_g \equiv GM/c^2$. The highest possible precession frequency occurs at the NS surface ($r = R$), giving $\nu_{LT} < 2\beta\nu_{\rm spin}(R_g/R)$. The maximum value for I is reached when a hollow sphere is assumed ($\beta = 2/3$). This is, of course, not a realistic assumption for a NS, we employ it merely as a hard upper limit for β. The true nature of the space-time external to such a shell is therefore irrelevant to our estimate. Assuming the NS radius to coincide with its own innermost stable circular orbit, which for such low spin is $r_{\rm ISCO} = 6R_g$ to a very good approximation, we find $\nu_{\rm LT} < 2.44$Hz. Higher order terms in a Kerr metric (e.g., Merloni et al. 1999), deviations of the metric from Kerr related to the NS structure (e.g., Morsink & Stella 1999), and arbitrarily inclined orbits (Sibgatullin 2002, e.g.,) could affect this upper limit by a factor of a few at most, and some of these effects actually lower it.

Considering that the entire inner disk precesses as a solid body between an inner and outer radius (e.g., Ingram & Done 2010) does not solve the discrepancy either, as in their description the entire disk cannot precess faster than a test particle at the inner radius. Our conclusion is that the 35–50 Hz QPO in IGR J17480–2446 can not be explained by frame dragging.

Figure 4 shows ν_ϕ, $\nu_{\rm peri}$ and $\nu_{\rm nodal}$ as a function of radius r for a point mass in the Kerr metric in an infinitesimally tilted and eccentric orbit (Merloni et al. 1999). In particular it shows a range of solutions for $\nu_{\rm nodal}$, with the upper limit resulting from assuming retrograde orbits and $R = r$. We note that theoretically the periastron precession frequency $\nu_{\rm peri}$ could in this system be identified with the QPOs at 35–50 Hz, but at a much larger radius (r ∼$20R_g$) than that where ν_ϕ is produced (r ∼$9R_g$). However, this would be entirely ad-hoc and would not work for the fast-spin NSs, where $\nu_{\rm peri}$ has instead been proposed to be identified with the lower kHz QPO.

Corrections due to classic precession are always of the order of, or lower than, $\nu_{\rm nodal}$ (under the assumption that deviations from a spherical NS are only due to the NS rotation, see Morsink & Stella 1999; see also Laarakkers & Poisson 1999). Shirakawa & Lai (2002) found that the NS magnetic field can induce warps in the inner accretion disk, resulting in a precessing inner flow. The resulting net precession frequency is set by a combination of (prograde) LT precession and (retrograde) classical and magnetic precession. The magnetic precession frequency $\nu_{\rm mag}$ is independent of $\nu_{\rm spin}$ and scales as μ^2, where μ is the magnetic moment. The likely higher μ in IGR J17480-2446 as compared with other NSs (Cavecchi et al. 2011; Papitto et al. 2011) suggests the possibility that magnetic precession could dominate in this system, opening a different possibility of explaining the LF-QPOs in IGR J17480–2446. Why in this scenario IGR J17480–2446 still conforms to the WK relation remains to be investigated, as $\nu_{\rm mag}$ depends on μ, the accretion rate, the viscosity of the disk, and the angle between the magnetic dipole and the spin axis. If magnetic precession does dominate in IGR J17480–2446, then magnetic effects would be expected to affect observed nodal precession frequencies in other, presumably lower magnetic-moment LMXBs as well.

In summary: the 35–50 Hz QPOs reported herein are incompatible with Lense-Thirring (or more generally GR nodal) precession, excluding frame dragging as the cause of the QPO in IGR J17480-2446. Given the similarities between the QPOs in IGR J17480-2446 and other Z-source systems, we conclude that frame dragging is in doubt as the mechanism that produces the horizontal-branch oscillations and, possibly, the hump component and LF-QPOs seen in Atoll sources and BH systems. While a scenario is conceivable where nodal precession causes all these QPOs, with magnetic precession dominating in IGR J17480–2446 and GR precession in the other systems, this requires a coincidence where systems with different magnetic fields all conform to a frequency-frequency relation (Figure 2) that previously was explained assuming no magnetic precession.

Acknowledgments: We are grateful to S. Morsink and P. Uttley for very insightful discussions. DA and MK acknowledge support from the International Space Science Institute (ISSI), Team Number 116. RW is partly supported by an ERC starting grant. J.H. acknowledges support from a NWO visitors grant.

REFERENCES

Altamirano D., van der Klis M., et al., 2008, ApJ, 685, 436
Altamirano D., Homan J., et al., Oct. 2010, The Astronomer's Telegram, 2952, 1
Altamirano et al. , Aug. 2012, To be submitted to ApJ., 0000, 1
Barret D., Jul. 2012, ApJ, 753, 84
Belloni T., Psaltis D., van der Klis M., 2002, ApJ, 572, 392
Belloni T.M., Sanna A., Mendez M., Jul. 2012, ArXiv e-prints
Casella P., Belloni T., Stella L., 2005, ApJ, 629, 403
Cavecchi Y., Patruno A., et al., Feb. 2011, ArXiv e-prints
Chakraborty M., Bhattacharyya S., Mukherjee A., Nov. 2011, MNRAS, 418, 490
Fragile P.C., Blaes O.M., et al., Oct. 2007, ApJ, 668, 417
Hasinger G., van der Klis M., 1989, A&A, 225, 79
Ingram A., Done C., Jul. 2010, MNRAS, 405, 2447
Ingram A., Done C., Aug. 2011, MNRAS, 415, 2323
Ingram A., Done C., Jan. 2012, MNRAS, 419, 2369
Ingram A., Done C., Fragile P.C., Jul. 2009, MNRAS, 397, L101
Jahoda K., Markwardt C.B., et al., 2006, ApJS, 163, 401
Jonker P.G., Wijnands R., et al., Jun. 1998, ApJ, 499, L191
Jonker P.G., van der Klis M., et al., Jul. 2000, ApJ, 537, 374
Klein-Wolt M., van der Klis M., Mar. 2008, ApJ, 675, 1407
Laarakkers W.G., Poisson E., Feb. 1999, ApJ, 512, 282
Leahy D.A., Darbro W., et al., 1983, ApJ, 266, 160
Lense J., Thirring H., 1918, Phys. Z. 19, 156
Linares M., van der Klis M., Wijnands R., 2007, ApJ, 660, 595
Linares M., Altamirano D., et al., Nov. 2011, ArXiv e-prints
Liu S., Melia F., Jul. 2002, ApJ, 573, L23
Méndez M., Belloni T., 2007, MNRAS, 381, 790
Merloni A., Vietri M., et al., Mar. 1999, MNRAS, 304, 155
Miyamoto S., Iga S., et al., 1993, ApJ, 403, L39
Morsink S.M., Stella L., Mar. 1999, ApJ, 513, 827
Olive J.F., Barret D., et al., 1998, A&A, 333, 942
Papitto A., D'Aì A., et al., Feb. 2011, A&A, 526, L3+
Patruno A., Watts A.L., Jun. 2012, ArXiv e-prints
Patruno A., Alpar M.A., et al., Jun. 2012, ApJ, 752, 33
Psaltis D., Belloni T., van der Klis M., 1999a, ApJ, 520, 262
Psaltis D., Wijnands R., et al., Aug. 1999b, ApJ, 520, 763
Remillard R.A., McClintock J.E., 2006, ARA&A, 44, 49
Rezzolla L., Yoshida S., et al., Sep. 2003, MNRAS, 344, L37
Shirakawa A., Lai D., Jan. 2002, ApJ, 564, 361
Sibgatullin N.R., Feb. 2002, Astronomy Letters, 28, 83
Stella L., Vietri M., 1998, ApJ, 492, L59+
Stella L., Vietri M., Morsink S.M., Oct. 1999, ApJ, 524, L63
Strohmayer T.E., Markwardt C.B., Oct. 2010, The Astronomer's Telegram, 2929, 1
van der Klis M., Jun. 1994, ApJS, 92, 511

van der Klis M., 1995, Proceedings of the NATO Advanced Study Institute on the Lives of the Neutron Stars, held in Kemer, Turkey, August 19-September 12, 1993. Editor(s), M. A. Alpar, U. Kiziloglu, J. van Paradijs; Publisher, Kluwer Academic, Dordrecht, The Netherlands, Boston, Massachusetts, 301

van der Klis M., 2006, in Compact Stellar X-Ray Sources, ed. W. H. G. Lewin & M. van der Klis (Cambridge: Cambridge Univ. Press)

van Straaten S., van der Klis M., Méndez M., 2003, ApJ, 596, 1155

Wagoner R.V., Silbergleit A.S., Ortega-Rodríguez M., Sep. 2001, ApJ, 559, L25

Warner B., Woudt P.A., 2002, MNRAS, 335, 84

Wijnands R., van der Klis M., 1999, ApJ, 514, 939

Zhang W., Jahoda K., et al., 1995, ApJ, 449, 930